

Generation of arbitrary two dimensional motional states of a trapped ion

XuBo Zou, K. Pahlke and W. Mathis

Institute TET, University of Hannover,
Appelstr. 9A, 30167 Hannover, Germany

Abstract

We present a scheme to generate an arbitrary two-dimensional quantum state of motion of a trapped ion. This proposal is based on a sequence of laser pulses, which are tuned appropriately to control transitions on the sidebands of two modes of vibration. Not more than $(M + 1)(N + 1)$ laser pulses are needed to generate a pure state with a phonon number limit M and N .

PACS number:42.50.Dv, 42.50.Ct

The generation of nonclassical states was studied in the past theoretically and experimentally. The first significant advances were made in quantum optics by demonstrating antibunched light [1] and squeezed light [2]. Various optical schemes for generating Schrödinger cat states were studied [3], which led to an experimental realization in a quantized cavity field [4]. Several schemes were proposed to generate any single-mode quantum state of a cavity field [5, 6, 7] and traveling laser field [8]. Recently, possible ways of generating various two-mode entangled field states were proposed. For example, it was shown [9] that entangled coherent states, which can be a superposition of two-mode coherent states [9, 10] can be produced using the nonlinear Mach-Zehnder interferometer. These quantum states can be considered as a two(multi)-mode generalization of single-mode Schrödinger cat states [11]. A method to generate another type of two-mode Schrödinger cat states, which are known as SU(2) Schrödinger cat states, was proposed [13, 14]. These quantum states result when two different SU(2) coherent states [12, 13, 14] are superposed. It was also shown that two-mode entangled number states can be generated by using nonlinear optical interactions [15], which may then be used to obtain the maximum sensitivity in phase measurements set by the Heisenberg limit [16]. In general, however, an experimental realization of nonclassical field states is difficult, because the quantum coherence can be destroyed easily by the interaction with the environment.

Recent advances in ion cooling and trapping have opened new prospects in nonclassical state generation. An ion confined in an electromagnetic trap can be described approximately as a particle in a harmonic potential. Its center of mass (c.m.) exhibits

a simple quantum-mechanical harmonic motion. By driving the ion appropriately with laser fields, its internal and external degrees of freedom can be coupled to the extent that its center-of-mass motion can be manipulated precisely. One advantage of the trapped ion system is that decoherence effects are relatively weak due to the extremely weak coupling between the vibrational modes and the environment. It was realized that this advantage of the trapped ion system makes it a promising candidate for constructing quantum logic gates for quantum computation [17] as well as for producing nonclassical states of the center-of-mass motion. In fact, single-mode nonclassical states, such as Fock states, squeezed states and Schrödinger cat states of the ion's vibration mode were investigated [18, 19, 20]. Recently, various schemes of producing two-mode nonclassical states of the vibration mode were proposed using ions in a two-dimensional trap [19, 21, 22]. In particular, several schemes were proposed to generate arbitrary two-dimensional motional quantum states of a trapped ion

$$\sum_{m=0}^M \sum_{n=0}^N C_{m,n} |m \rangle_x |n \rangle_y . \quad (1)$$

The ket-vectors $|m \rangle_x$ and $|n \rangle_y$ denote the Fock states of the quantized vibration along the x and y axes. In the schemes of Gardiner et al [23], the number of required laser operations depends exponentially on the upper photon numbers M and N . The scheme introduced by Kneer and Law [24] involves $(2M+1)(N+1)+2N$ operations, while the scheme proposed by Drobny et al [25] requires $2(M+N)^2$ operations. More recently S. B. Zheng presented a scheme [26], which requires only $(M+2)(N+1)$ operations. The reduction of the number of operation is important for further experimental realization. Notice that the quantum state (1) involves $(M+1)(N+1)$ complex coefficients. The purpose of this paper is to present a scheme to generate quantum state (1) with not more than $(M+1)(N+1)$ quantum operations. We will show that each coefficient of quantum state (1) corresponds to only one laser pulse of this most efficient quantum state generation scheme. In order to describe this concept in an obvious way, we split the Hilbert space H_{vib} of the two modes of vibration to subspaces H_{vib}^{2J} with a constant total number of vibrational quanta $m+n = 2J$. Thus, the Hilbert-space is a direct sum: $H_{vib} = \bigcup_{J=0}^{\infty} H_{vib}^{2J}$ with $2J \in \mathbb{N}$. The quantum states are formulated by the two-mode Fock states $|J, L \rangle = |J+L \rangle_x |J-L \rangle_y$ with $L \in \{-J, -J+1, \dots, J\}$. So J and L can be half numbers. The subspaces H_{vib}^{2J} are spanned by $|J, L \rangle; |J, L-1 \rangle; \dots; |J, -L \rangle$. The quantum state (1) can also be written in the form

$$\sum_{J=0}^{(M+N)/2} \sum_{L=-J}^J d_{J,L} |J, L \rangle . \quad (2)$$

In order to synthesize 2D quantum states of the vibration of one trapped ion we suggest to use laser stimulated Raman processes [27]. We consider a trapped ion confined in a 2D harmonic potential characterized by the trap frequencies ν_x and ν_y in two orthogonal directions x and y . The ion is irradiated along the x and y axes by two external laser frequencies ω_x, ω_y and wave vectors k_x, k_y . The two laser fields stimulate Raman transitions between two electronic ground state levels $|g_1 \rangle$ and $|g_2 \rangle$ via a third electronic level, which is far enough out of resonance. The difference of energy of these two electronic ground states is $\hbar\omega_0$. We concentrate our investigations on those

transitions on the quantum states of the trapped ion, which are in resonance with the laser field

$$\omega_x - \omega_y = \omega_0 - m\nu_x - n\nu_y; \quad m, n \in \mathbb{N} . \quad (3)$$

This equation can be interpreted as a condition of resonance, which relates the difference of frequency of two Raman laser beams to quantum numbers of two modes of vibration. The concept of the quantum state manipulation scheme, which we present, is based on appropriately tuned laser pulses to drive sideband transitions with different numbers n, m selectively. Thus, we want to control multi-phonon transitions with simultaneous changes of the vibrational motion in both directions. We intend to generate each component of the quantum state (1) from the ground state of the collective vibration. To make this concept reliable we assume incommensurate trap frequencies ν_x and ν_y . This requirement can be fulfilled by trap design, since the number of photons in the quantum state (1) is bounded: $m \leq M, n \leq N$.

If the resonance condition (3) holds for every laser pulse the modeling can be simplified. In the example of a single ion confined in a two-dimensional harmonic trap Steinbach [27] set a slightly different condition to describe the phonon exchange between different directions of vibrational motion. We do a similar kind of modelling, but with a different goal. After the standard dipole and rotating wave approximation, the adiabatic elimination of the auxiliary off-resonant levels lead to an effective Hamiltonian

$$H = H_0 + H_{int} \quad (4)$$

with

$$H_0 = \frac{\omega_0}{2}(|g_2 \rangle \langle g_2| - |g_1 \rangle \langle g_1|) + \nu_x a^\dagger a + \nu_y b^\dagger b; \quad (5)$$

$$H_{int} = \Omega e^{i\eta_x(a+a^\dagger)+i\eta_y(b+b^\dagger)+i(\omega_x-\omega_y)t} |g_2 \rangle \langle g_1| + h.c.. \quad (6)$$

Where a and b ($a^\dagger$ and $b^\dagger$) are annihilation(creation) operators of the quantized motion along the x and y axes. η_x and η_y are the associated Lamb-Dicke parameters in the x and y direction. The complex parameter Ω denotes the effective Raman coupling constant, which considers the phase difference ϕ of the two Raman laser beams.

In order to generate the motional quantum state (2), we consider the situation in which the ion is prepared in the ground state $|g_2 \rangle$ and the two c.m. modes in the vacuum states $|0 \rangle_x$ and $|0 \rangle_y$:

$$\Psi_{initial} = |g_2 \rangle |0 \rangle_x |0 \rangle_y = |g_2, 0, 0 \rangle . \quad (7)$$

In the following we show, that each term of this linear combination can be generated from the ground state term $|0 \rangle_x |0 \rangle_y$ most efficiently by one laser pulse.

In the rotating wave approximation we consider only those terms of the operator H_{int} , which are in resonance with the actual laser pulse. That is why we write the operator of interaction conveniently in the form

$$\begin{aligned} H_{m,n} = & \Omega e^{-\eta_x^2/2 - \eta_y^2/2} \sum_{k_1, k_2=0}^{\infty} \frac{(i\eta_x)^{2k_1+m} (i\eta_y)^{2k_2+n}}{k_1! k_2! (k_1+m)! (k_2+n)!} a^{\dagger k_1} a^{k_1} b^{\dagger k_2} b^{k_2} a^m b^n |g_2 \rangle \langle g_1| \\ & + h.c.. \end{aligned} \quad (8)$$

The index $\{m, n\}$ indicates the relevant components of H_{int} , which correspond to the condition of resonance (3). It follows, that the operator of time evolution $U_{mn} = \exp(-iH_{m,n}t)$ satisfies the relations

$$U_{mn}|g_2 > |0 >_x |0 >_y = \cos(|\Omega_{m,n}|t)|g_2 > |0 >_x |0 >_y - ie^{-i\phi_{m,n}} \sin(|\Omega_{m,n}|t)|g_1 > |m >_x |n >_y \quad (9)$$

$$U_{mn}|g_1 > |k >_x |l >_y = |g_1 > |k >_x |l >_y; \quad k + l < m + n \quad (10)$$

$$U_{mn}|g_1 > |k >_x |l >_y = |g_1 > |k >_x |l >_y; \quad k + l = m + n; k \neq m. \quad (11)$$

Here $|\Omega_{m,n}|$ and $\phi_{m,n}$ are the amplitude and the phase of the corresponding Raman coupling constant

$$\Omega_{m,n} = |\Omega_{m,n}| e^{i\phi_{m,n}} = \Omega e^{-\eta_x^2/2 - \eta_y^2/2 + i\phi} \frac{(i\eta_x)^m (i\eta_y)^n}{m!n!}. \quad (12)$$

We begin from the ground state (7) by applying the first laser pulse. It fulfills the resonance condition (3) in the case of: $n = 0; m = 0$. This laser pulse, which corresponds to $H_{0,0}$, has to generate the term $d_{0,0}|0, 0 >$ of the quantum state (2). We choose the duration $t_{0,0}$ of the laser pulse to fulfill

$$-ie^{-i\phi_{0,0}} \sin(|\Omega_{0,0}|t_{0,0}) = d_{0,0}. \quad (13)$$

The system's state is transformed into

$$\Psi^{0,0} = \cos(|\Omega_{0,0}|t_{0,0})|g_2 > |0, 0 > -ie^{-i\phi_{0,0}} \sin(|\Omega_{0,0}|t_{0,0})|g_1 > |0, 0 > \quad (14)$$

$$= \sqrt{1 - |d_{0,0}|^2}|g_2 > |0, 0 > + d_{0,0}|g_1 > |0, 0 >. \quad (15)$$

In order to drive the system with the effective interaction $H_{0,1}$ the next laser pulse is tuned according to Eq. (3) to resonance: $(m; n) = (0; 1)$. After this laser pulse has driven the ion with a duration $t_{0,1}$, the system's state becomes

$$\begin{aligned} \Psi^{0,1} &= \sqrt{1 - |d_{0,0}|^2} [\cos(|\Omega_{0,1}|t_{0,1})|g_2 > |0, 0 > \\ &\quad - ie^{-i\phi_{0,1}} \sin(|\Omega_{0,1}|t_{0,1})|g_1 > |\frac{1}{2}, -\frac{1}{2} >] + d_{0,0}|g_1 > |0, 0 >. \end{aligned} \quad (16)$$

We choose the laser pulse duration $t_{0,1}$ to satisfy

$$-ie^{-i\phi_{0,1}} \sqrt{1 - |d_{0,0}|^2} \sin(|\Omega_{0,1}|t_{0,1}) = d_{\frac{1}{2}, -\frac{1}{2}}. \quad (17)$$

Thus, after the $(\frac{1}{2}, -\frac{1}{2})$ -component of Eq. (2) is generated, the system's state becomes

$$\begin{aligned} \Psi^{0,1} &= \sqrt{1 - |d_{0,0}|^2 - |d_{\frac{1}{2}, -\frac{1}{2}}|^2} |g_2 > |0, 0 > \\ &\quad + d_{\frac{1}{2}, -\frac{1}{2}} |g_1 > |\frac{1}{2}, -\frac{1}{2} > + d_{0,0}|g_1 > |0, 0 >. \end{aligned} \quad (18)$$

We proceed with a laser pulse, which is characterized by $(m; n) = (1; 0)$, to let the quantum state evolve into

$$\begin{aligned}\Psi^{1,0} = & \sqrt{1 - |d_{0,0}|^2 - |d_{\frac{1}{2}, -\frac{1}{2}}|^2} [\cos(|\Omega_{1,0}|t_{1,0})|g_2 > |0, 0 > \\ & -ie^{-i\phi_{1,0}} \sin(|\Omega_{1,0}|t_{1,0})|g_1 > |\frac{1}{2}, \frac{1}{2} >] \\ & + d_{\frac{1}{2}, -\frac{1}{2}}|g_1 > |\frac{1}{2}, -\frac{1}{2} > + d_{0,0}|g_1 > |0, 0 > .\end{aligned}\quad (19)$$

With the choice

$$-ie^{-i\phi_{1,0}} \sqrt{1 - |d_{0,0}|^2 - |d_{\frac{1}{2}, -\frac{1}{2}}|^2} \sin(|\Omega_{1,0}|t_{1,0}) = d_{\frac{1}{2}, \frac{1}{2}} \quad (20)$$

we obtain

$$\begin{aligned}\Psi^{1,0} = & \sqrt{1 - |d_{0,0}|^2 - |d_{\frac{1}{2}, -\frac{1}{2}}|^2 - |d_{\frac{1}{2}, \frac{1}{2}}|^2} |g_2 > |0, 0 > \\ & + d_{\frac{1}{2}, \frac{1}{2}}|g_1 > |\frac{1}{2}, \frac{1}{2} > + d_{\frac{1}{2}, -\frac{1}{2}}|g_1 > |\frac{1}{2}, -\frac{1}{2} > + d_{0,0}|g_1 > |0, 0 > .\end{aligned}\quad (21)$$

If this procedure is done for the $[\frac{(i+j)(i+j+1)}{2} + i + 1]$ -th time, the quantum state of the system is

$$\begin{aligned}\Psi^{i,j} = & \sum_{J=0}^{(i+j-1)/2} \sum_{L=-J}^J d_{J,L} |J, L > |g_1 > + \sum_{L=-(i+j)/2}^{(i-j)/2} d_{(i+j)/2, L} |(i+j)/2, L > |g_1 > \\ & + \sqrt{1 - \sum_{J=0}^{(i+j-1)/2} \sum_{L=-J}^J |d_{J,L}|^2 - \sum_{L=-(i+j)/2}^{(i-j)/2} |d_{(i+j)/2, L}|^2} |g_2 > |0, 0 > .\end{aligned}\quad (22)$$

We now consider the $[\frac{(i+j)(i+j+1)}{2} + i + 2]$ -th operation by discussing the two possible cases.

In the special case of $j = 0$ we drive the ion with the operator of interaction $H_{0,i+1}$. After the system has been driven by the corresponding laser pulse of duration $t_{0,i+1}$, the quantum state becomes

$$\begin{aligned}\Psi^{0,i+1} = & \sum_{J=0}^{(i)/2} \sum_{L=-J}^J d_{J,L} |J, L > |g_1 > \\ & + \sqrt{1 - \sum_{J=0}^{(i)/2} \sum_{L=-J}^J |d_{J,L}|^2} [\cos(|\Omega_{0,i+1}|t_{0,i+1})|g_2 > |0, 0 > \\ & -ie^{-i\phi_{0,i+1}} \sin(|\Omega_{0,i+1}|t_{0,i+1})|g_1 > |(i+1)/2, -(i+1)/2 >].\end{aligned}\quad (23)$$

With the choice

$$-ie^{-i\phi_{0,i+1}} \sqrt{1 - \sum_{J=0}^{(i)/2} \sum_{L=-J}^J |d_{J,L}|^2} \sin(|\Omega_{0,i+1}|t_{0,i+1}) = d_{(i+1)/2, -(i+1)/2} \quad (24)$$

we obtain

$$\begin{aligned}\Psi^{0,i+1} &= \sum_{J=0}^{(i)/2} \sum_{L=-J}^J d_{J,L} |J, L\rangle |g\rangle + d_{(i+1)/2, -(i+1)/2} |(i+1)/2, -(i+1)/2\rangle |g_1\rangle \\ &+ \sqrt{1 - \sum_{J=0}^{(i)/2} \sum_{L=-J}^J |d_{J,L}|^2 - |d_{(i+1)/2, -(i+1)/2}|^2} |g_2\rangle |0, 0\rangle .\end{aligned}\quad (25)$$

In the other case ($j \neq 0$) the ion is driven by a laser pulse, which corresponds to $H_{i+1,j-1}$. We choose the interaction time $t_{i+1,j-1}$ and phase of laser field to satisfy

$$\begin{aligned}d_{(i+j)/2, (i-j+2)/2} &= -ie^{-i\phi_{i+1,j-1}} \sin(|\Omega_{i+1,j-1}| t_{i+1,j-1}) \\ &\times \sqrt{1 - \sum_{J=0}^{(i+j-1)/2} \sum_{L=-J}^J |d_{J,L}|^2 - \sum_{L=-(i+j)/2}^{(i-j)/2} |d_{(i+j)/2, L}|^2}\end{aligned}\quad (26)$$

and to obtain the quantum state

$$\begin{aligned}\Psi^{i+1,j-1} &= \sum_{J=0}^{(i+j-1)/2} \sum_{L=-J}^J d_{J,L} |J, L\rangle |g_1\rangle + \sum_{L=-(i+j)/2}^{(i-j+2)/2} d_{(i+j)/2, L} |(i+j)/2, L\rangle |g_1\rangle \\ &+ \sqrt{1 - \sum_{J=0}^{(i+j-1)/2} \sum_{L=-J}^J |d_{J,L}|^2 - \sum_{L=-(i+j)/2}^{(i-j+2)/2} |d_{(i+j)/2, L}|^2} |g_2\rangle |0, 0\rangle .\end{aligned}\quad (27)$$

After the procedure is performed for $\frac{(M+N+1)(M+N)}{2}$ times the system's state definitely becomes

$$\Psi_{final} = \sum_{J=0}^{(M+N)/2} \sum_{L=-J}^J d_{J,L} |J, L\rangle |g_1\rangle .\quad (28)$$

Thus, the system is prepared in a product state Ψ of the desired vibrational quantum state (2) and the ground state $|g_1\rangle$.

We have proposed a scheme to generate any two-dimensional quantum state of vibration of one trapped ion. This concept is based on the possibility to drive sideband transitions with different numbers n, m selectively by tuning laser pulses appropriately. Thus, we want to control multi-phonon transitions with simultaneous changes of the quantum numbers of vibration in both directions. Each component of the quantum state (1) can be generated step by step from the ground state of vibration. To make our concept reliable we have assumed that the trap frequencies ν_x and ν_y can be made incommensurate by trap design.

However, there might exist on-resonant terms in addition to the ones included in Eq. (8) due to finite bandwidth of laser pulse. If the ratio of the trap frequencies is chosen appropriately so that frequency of each laser pulse is enough separated, these terms can be neglected in the Lamb-Dicke limit. We assume equal values of the Lamb-Dicke parameter $\eta_x = \eta_y$ [27] and require a ratio of trap frequencies to satisfy condition

$\nu_x/\nu_y > M + 2N$. In this case, we see that the laser frequency of each sideband transition is enough separated and we can neglect those unwanted resonant terms. Thus, in order to implement our scheme, we require a large enough trap anisotropies. In addition, in Lamb-Dicke limit, the effective Rabi frequency $\Omega_{m,n}$ of Eq. (12) is proportional to $\frac{(i\eta_x)^m(i\eta_y)^n}{m!n!}$. If the quantum numbers of vibration n and m are too large the effective Rabi frequency $\Omega_{m,n}$ becomes too small. Thus, in respect of decoherence the time required to complete the procedure might become too long. This problem represents the limitation of this scheme.

This scheme of quantum state generation requires not more than $(M+1)(N+1)$ laser pulses, which is the smallest possible number. Thus, in respect of short laser pulse sequences it can be considered as the optimal solution of this problem of the quantum state generation. The number of required laser pulses reduces, if there are coefficients in the target quantum state, which are equal to zero. For example, if the coefficient $d_{J,L}$ of Eq.(2), which is generated with the $(2J^2 + J + L + 1)$ -th operation, is zero, it is not necessary to apply the corresponding laser pulse.

References

- [1] See, for a review, H. Paul, Rev. Mod. Phys. **54**, 1061 (1982); R. Loudon, Rep. Prog. Phys. **43**, 913 (1980).
- [2] See, for a review, R. Loudon and P. L. Knight, J. Mod. Opt. **34**, 709 (1987).
- [3] B. Yurke and D. Stoler, Phys. Rev. Lett. **57**, 13 (1986); B. Yurke and D. Stoler, Phys. Rev. A **35**, 4846 (1987); M. Brune, S. Haroche, J. M. Raimond, L. Davidovich, and N. Zagury, Phys. Rev. A **45**, 5193 (1992).
- [4] M. Brune, E. Hagley, J. Dreyer, X. Maitre, A. Maali, C. Wunderlich, J. M. Raimond, and S. Haroche, Phys. Rev. Lett. **77**, 4887 (1996).
- [5] K. Vogel, V. M. Akulin and W. P. Schleich, Phys. Rev. Lett. **71**, 1816 (1993).
- [6] A. S. Parkins, P. Marte, P. Zoller and H. J. Kimble, Phys. Rev. Lett. **71**, 3095 (1993).
- [7] C. K. Law and J. H. Eberly, Phys. Rev. Lett. **76**, 1055 (1996).
- [8] M. Dakna, J. Clausen, L. Knöll and D. G. Welsch, Phys. Rev. A **59**, 1658 (1999).
- [9] B. C. Sanders, Phys. Rev. A **45**, 6811 (1992); B. C. Sanders and D. A. Rice, Phys. Rev. A **61**, 013805 (1999).
- [10] C. L. Chai, Phys. Rev. A **46**, 7187 (1992).
- [11] N. A. Ansari and V. I. Man'ko, Phys. Rev. A **50**, 1942 (1994); V. V. Dodonov, V. I. Man'ko, and D. E. Nikonov, Phys. Rev. A **51**, 3328 (1995).
- [12] V. Buzek and T. Quang, J. Opt. Soc. Amer. B **6**, 2447 (1989).

- [13] B. C. Sanders, Phys. Rev. A **40**, 2417 (1989).
- [14] C. C. Gerry and R. Grobe, J. Mod. Opt. **44**, 41 (1997).
- [15] C. C. Gerry, Phys. Rev. A **61**, 043811 (2000).
- [16] J. J. Bollinger, W. M. Itano, D. J. Wineland, and D. J. Heinzen, Phys. Rev. A **54**, R4649 (1996).
- [17] J. I. Cirac and P. Zoller, Phys. Rev. Lett. **74**, 4091 (1995).
- [18] J. I. Cirac, R. Blatt, A. S. Parkins, and P. Zoller, Phys. Rev. Lett. **70**, 762 (1993);
J. I. Cirac, A. S. Parkins, R. Blatt, and P. Zoller, Phys. Rev. Lett. **70**, 556 (1993);
R. L. de Matos Filho and W. Vogel, Phys. Rev. Lett. **76**, 608 (1996); J. F. Poyatos,
J. I. Cirac, R. Blatt, and P. Zoller, Phys. Rev. A **54**, 1532 (1996); S. B. Zheng,
Phys. Rev. A **58**, 761 (1998).
- [19] C. C. Gerry, Phys. Rev. A **55**, 2478 (1997).
- [20] D. M. Meekhof, C. Monroe, B. E. King, W. M. Itano, and D. J. Wineland, Phys.
Rev. Lett. **76**, 1796 (1996); C. Monroe, D. M. Meekhof, B. E. King, and D. J.
Wineland, Science **272**, 1131 (1996).
- [21] S. C. Gou and P. L. Knight, Phys. Rev. A **54**, 1682 (1996).
- [22] S. C. Gou, J. Steinbach, and P. L. Knight, Phys. Rev. A **54**, R1014 (1996); S. C.
Gou, J. Steinbach, and P. L. Knight, Phys. Rev. A **54**, 4315 (1996); S. Maniscalco,
A. Messina, and A. Napoli, Phys. Rev. A **61**, 053806 (2000). X. B. Zou, J. W.
Kim and H. W. Lee, Phys. Rev. A **63**, 065801 (2001)
- [23] S. A. Gardiner, J. I. Cirac and P. Zoller, Phys. Rev. A **55**, 1683 (1997).
- [24] B. Kneer and C. K. Law, Phys. Rev. A **57**, 2096 (1998).
- [25] G. Dronby, B. Hladky and V. Buzek, Phys. Rev. A **58**, 2481 (1995).
- [26] S. B. Zheng, Phys. Rev. A **63**, 015801 (2000).
- [27] J. Steinbach, J. Twamley, and P. L. Knight, Phys. Rev. A **56**, 4815 (1997).